

Краят на диетите

Д-р Джоуел Фърмън

Издаелство Еуниката
София, 2016

Бележки и библиография

Обетът за край на диетите

1. Franz MJ, Van Wormer JJ, Crain AL, et al. Weight-loss outcomes: a systematic review and meta-analysis of weight-loss clinical trials with a minimum 1-year follow-up. *J Am Diet Assoc.* 2007;107(10):1755–1767.
2. Montani JP, Viecelli AK, Prévot A, Dulloo AG. Weight cycling during growth and beyond as a risk factor for later cardiovascular diseases: the “repeated overshoot” theory. *Int J Obes (Lond).* 2006;30(suppl 4):S58–66.

Увод

1. Mokdad AH, Marks JS, Stroup DF, Gerberding JL. Actual causes of death in the United States, 2000. *JAMA.* 2004;291(10):1238–1245.

Първа глава: Токсичен глад

1. Lennerz BS, Alsop DC, Holsen LM, et al. Effects of dietary glycemic index on brain regions related to reward and craving in men. *Am J Clin Nutr.* 2013; 98(3):641–7.
2. Science Daily. Compulsive eating shares addictive biochemical mechanism with cocaine, heroin abuse, study shows. <http://www.sciencedaily.com/releases/2010/03/100328170243.htm>. Published March 29, 2010. Accessed April 16, 2010.
3. Yona MA, Maugera SL, Pickavance LC. Relationships between dietary macronutrients and adult neurogenesis in the regulation of energy metabolism. *Brit J Nutrition.* 2013;109(9):1573–1589.
4. Holt-Lunstad J, Smith TB, Layton JB. Social relationships and mortality risk: a meta-analytic review. *PLoS Med.* 2010;7(7):e1000316. doi: 10.1371/journal.pmed.1000316.

Втора глава: Митовете за храненето развенчани

1. Biba E. Where is the next Carl Sagan? *Popular Science.* June 3, 2013, <http://www.popsci.com/science/article/2013-05/not-just-facts>.
2. Sacks FM, Bray GA, Carey VJ, et al. Comparison of weight-loss diets with different compositions of fat, protein, and carbohydrates. *N Engl J Med.* 2009;360(9):859–873. De Souza RJ, Bray GA, Carey VJ, et al. Effects of 4 weight-loss diets differing in fat, protein, and carbohydrate on fat mass, lean mass, visceral adipose tissue, and hepatic fat: results from the

- POUNDS LOST trial. *Am J Clin Nutr.* 2012;95(3):614–625. Dansinger ML, Gleason JA, Griffith JL, et al. Comparison of the Atkins, Ornish, Weight Watchers, and Zone diets for weight loss and heart disease risk reduction: a randomized trial. *JAMA.* 2005;293(1):43–53.
3. Estruch R, Ros E, Salas-Salvadó J, et al. Primary prevention of cardiovascular disease with a Mediterranean diet. *N Engl J Med.* 2013;368:1279–1290.
 4. Note to come.
 5. Kramer CK, Zinman B, Retnakaran R. Are metabolically healthy overweight and obesity benign conditions? A systemic review and meta-analysis. *Ann Intern Med.* 2013;159(11):758–69.
 6. Flegal KM, Kit BK, Orpana H, et al. Association of all-cause mortality with overweight and obesity using standard body mass index categories: a systemic review and meta-analysis. *JAMA.* 2013;309:71–82.
 7. Singh PN, Fraser GE. Dietary risk factors for colon cancer in a low-risk population. *Am J Epidemiol.* 1998;148(8):761–774. Aune D, De Stefani E, Ronco A, et al. Legume intake and the risk of cancer: a multisite case-control study in Uruguay. *Cancer Causes Control.* 2009;20(9):1605–1615. Darmadi-Blackberry I, Wahlqvist ML, Kouris-Blazos A, et al. Legumes: the most important dietary predictor of survival in older people of different ethnicities. *Asia Pac J Clin Nutr.* 2004;13(2):217–220.
 8. Chitnis MM, Yuen JS, Protheroe AS, et al. The type 1 insulin-like growth factor receptor pathway. *Clin Cancer Res.* 2008;14:6364–6370. Werner H, Bruchim I. The insulin-like growth factor-I receptor as an oncogene. *Arch Physiol Biochem.* 2009;115:58–71. Davies M, Gupta S, Goldspink G, et al: The insulin-like growth factor system and colorectal cancer: clinical and experimental evidence. *Int J Colorectal Dis.* 2006;21:201–208. Sandhu MS, Dunger DB, Giovannucci EL. Insulin, insulin-like growth factor-I (IGF-I), IGF binding proteins, their biologic interactions, and colorectal cancer. *J Natl Cancer Inst.* 2002;94:972–980. Rowlands MA, Gunnell D, Harris R, Vatten LJ, Holly JM, Martin RM. Circulating insulin-like growth factor peptides and prostate cancer risk: a systematic review and meta-analysis. *Int J Cancer.* 2009 May 15;124(10):2416–29.
 9. Bartke A. Minireview: role of the growth hormone/insulin-like growth factor system in mammalian aging. *Endocrinology.* 2005;146:3718–23.
 10. Fung TT, van Dam RM, Harkinson SE, et al. Low-carbohydrate diets and all-cause and cause-specific mortality: two cohort studies. *Ann Intern Med.* 2010;153:289–298.
 11. Preis SR, Stampfer MJ, Spiegelman D, et al. Dietary protein and risk of ischemic heart disease in middle-aged men. *Am J Clin Nutr.* 2010;92:1265–1272.

12. Ligiou P, Sandin S, Lof M, et al. Low carbohydrate, high protein diet and incidence of cardiovascular disease in Swedish women: prospective cohort study. *BMJ*. 2012;344:e4026. doi: 10.1136/bmj.e4026.
13. Pan A, Sun Q, Bernstein AM, et al. Changes in red meat consumption and subsequent risk of type 2 diabetes mellitus: three cohorts of US men and women. *JAMA Intern Med*. 2013;173(14):1328–1335.
14. Vergnaud AC, Norat T, Romaguera D, et al. Meat consumption and prospective weight change in participants of the EPIC-PANACEA study. *Am J Clin Nutr*. 2010;92(2):398–407.
15. Fraser GE, Shavlik DJ. Ten years of life: is it a matter of choice? *Arch Intern Med*. 2001;161:1645–1652.
16. Guasch-Ferre M, Bulio M, Gartinez-Gonzalez MA, et al. Frequency of nut consumption and mortality risk in the PREDIMED nutrition intervention trial. *BMC Med*. 2013;11:164. doi:10.1186/1741-7015-11-164. Rohrmann S, Faeh D. Should we go nuts about nuts? *BMC Med*. 2013;11:165. doi:10.1186/1741-7015-11-165.
17. Sofi F, Vecchio S, Giuliani G, et al. Dietary habits, lifestyle and cardiovascular risk factors in a clinically healthy Italian population: the 'Florence' diet is not Mediterranean. *European Journal of Clinical Nutrition*. 2005;59, 584–591.
18. Salas-Salvadó J, Casas-Agustench P, Murphy MM, et al. The effect of nuts on inflammation. *Asia Pac J Clin Nutr*. 2008;17(suppl 1):333–336. Casas-Agustench P, Bulló M, Salas-Salvadó J. Nuts, inflammation and insulin resistance. *Asia Pac J Clin Nutr*. 2010;19(1):124–130. Gopinath B, Buyken AE, Flood VM, et al. Consumption of polyunsaturated fatty acids, fish, and nuts and risk of inflammatory disease mortality. *Am J Clin Nutr*. 2011;93(5):1073–1079. Ros E. Nuts and novel biomarkers of cardiovascular disease. *Am J Clin Nutr*. 2009;89(5):1649S–1656S. Kris-Etherton PM, Hu FB, Ros E, Sabaté J. The role of tree nuts and peanuts in the prevention of coronary heart disease: multiple potential mechanisms. *J Nutr*. 2008;138(9):1746S–1751S. O'Neil CE, Keast DR, Nicklas TA, Fulgoni VL III. Nut consumption is associated with decreased health risk factors for cardiovascular disease and metabolic syndrome in U.S. adults: NHANES 1999–2004. *J Am Coll Nutr*. 2011;30(6):502–510. Falasca M, Casari I. Cancer chemoprevention by nuts: evidence and promises. *Front Biosci (Schol Ed)*. 2012;1(4):109–120. González CA, Salas-Salvadó J. The potential of nuts in the prevention of cancer. *Br J Nutr*. 2006;96(suppl 2):S87–94.
19. Sala-Vila A, Romero-Mamani ES, Gilibert R, et al. Mediterranean diet and 2.4-year changes in ultrasound-assessed carotid intima-media thickness and plaque: a sub-study of the PREDIMED trial. Presented at the 81st EAS Congress, June 2–5, 2013, Lyons, France.

20. Guasch-Ferré M, Bulló M, Martínez-González MA, et al. Frequency of nut consumption and mortality risk in the PREDIMED nutrition intervention trial. *BMC Med.* 2013;11:164.
21. Rubio-Tapia A, Ludvigsson JF, Brantner TL, et al. The prevalence of celiac disease in the United States. *Am J Gastroenterol.* 2012;107(10):1538–44.
2. Sapone A, Bai JC, Ciacci C, et al. Spectrum of gluten-related disorders: consensus on new nomenclature and classification. *BMC Med.* 2012;10:13. National Foundation for Celiac Awareness: Celiac Disease & Non-Celiac Gluten Sensitivity. http://www.celiaccentral.org/SiteData/docs/NFCA_Celiac/a5c2249c6b6762ab/NFCA_CeliacDisease_vs_NonCeliacGlutenSensitivity.pdf.
22. Note TK.
23. Ornish D, Brown SE, Scherwitz LW, et al. Can lifestyle changes reverse coronary heart disease? The Lifestyle Heart Trial. *Lancet.* 1990;336(8708):129–133.
24. Esselstyn CB Jr. Updating a 12-year experience with arrest and reversal therapy for coronary heart disease (an overdue requiem for palliative cardiology). *Am J Cardiol.* 1999;84:339–341, A338.
25. Jaceldo-Siegl K, Oda K, St John C, et al. Nut intake and risk of metabolic syndrome. *FASEB J.* 2013;27:120.8. Kendall CWC, Augustin LSA, Bashyam B, et al. Effect of nuts on coronary heart disease risk factors in type 2 diabetes. *FASEB J.* 2013;27:368.5. O’Neil CE, Nicklas TA, Fulgoni VL. Tree nut consumption is associated with better diet quality, nutrient intake of select nutrients, and better measures of some cardiovascular risk factors (CVRF): National Health and Nutrition Examination Survey (NHANES) 2005–2010. *FASEB J.* 2013;27:847.13.
26. Baer HJ, Glynn RJ, Hu FB, et al. Risk factors for mortality in the nurses’ health study: a competing risks analysis. *Am J Epidemiol.* 2011;173(3):319–329.
27. Fraser GE, Shavlik DJ. Ten years of life: is it a matter of choice? *Arch Intern Med.* 2001;161:1645–1652.
28. Schaefer EJ, Bongard V, Beiser AS, et al. Plasma phosphatidylcholine docosahexaenoic acid content and risk of dementia in Alzheimer disease. *Arch Neurol.* 2006;63:1545–1550.
29. van Gelder BM, Tijhuis M, Kalmijn S, Kromhout D. Fish consumption, n-3 fatty acids, and subsequent 5-y cognitive decline in elderly men: the Zutphen Elderly Study. *Am J Clin Nutr.* 2007;85:1142–1147. Beydoun MA, Kaufman JS, Satia JA, et al. Plasma n-3 fatty acids and the risk of cognitive decline in older adults: the Atherosclerosis Risk in Communities Study. *Am J Clin Nutr.* 2007;85:1103–1011.

30. Söderberg M, Edlund C, Kristensson K, Dallner G. Fatty acid composition of brain phospholipids in aging and in Alzheimer's disease. *Lipids*. 1991;26:421–425.
31. Huang TL. Omega-3 fatty acids, cognitive decline, and Alzheimer's disease: a critical review and evaluation of the literature. *J Alzheimers Dis*. 2010;21:673–690.
32. Yurko-Mauro K, McCarthy D, Rom D, et al. Beneficial effects of docosahexaenoic acid on cognition in age-related cognitive decline. *Alzheimers Dement*. 2010;6:456–464.
33. Stonehouse W, Conlon CA, Podd J, et al. DHA supplementation improved both memory and reaction time in healthy young adults: a randomized controlled trial. *Am J Clin Nutr*. 2013;97(5):1134–1143.
34. Lewis MD, Hibbeln JR, Johnson JE, et al. Suicide deaths of active-duty US military and omega-3 fatty-acid status: a case-control comparison. *J Clin Psychiatry* 2011;72(12):1585–1590.
35. Sublette ME, Ellis SP, Geant AL, Mann JJ. Meta-analysis of the effects of eicosapentaenoic acid in clinical trials in depression. *J Clin Psychiatry*. 2011;72(12):1577–1584.
36. Sarter B, Kelsey K, Harris W. Blood docosahexaenoic acid and eicosapentaenoic acid in vegans: associations with age and gender and effects of supplementation. Personal communication, publication pending.
37. Geppert J, Kraft V, Demmelmair H, Koletzko B. Docosahexaenoic acid supplementation in vegetarians effectively increases omega-3 index: a randomized trial. *Lipids*. 2005;40(8):807–814.
38. Hu FB, Stampfer MJ. Nut consumption and risk of coronary heart disease: a review of epidemiologic evidence. *Curr Atheroscler Rep*. 1999;1(3):204–209.

Трета глава: Краят на диетите

1. Lauer JB, Reed GW, Hill JO. Effects of weight cycling induced by diet cycling in rats differing in susceptibility to dietary obesity. *Obes Res*. 1999;7: 215–222. Reed GW, Cox G, Yakubu F, Ding L, Hill JO. Effects of weight cycling in rats allowed a choice of diet. *Am J Physiol*. 1993;264:R35–R40.
2. Wallner SJ, Luschnigg N, Schnedl WJ, et al. Body fat distribution of overweight females with a history of weight cycling. *Int J Obes Relat Metab Disord*. 2004;28:1143–1148. Van der Kooy K, Leenen R, Seidell JC, et al. Effect of a weight cycle on visceral fat accumulation. *Am J Clin Nutr*. 1993;58:853–857.
3. Olson MB, Kelsey SF, Bittner V, et al. Weight cycling and high-density lipoprotein cholesterol in women: evidence of an adverse effect: a report from the NHLBI-sponsored WISE study. Women's Ischemia Syndrome

- Evaluation Study Group. *J Am Coll Cardiol*. 2000;36:1565–1571. Rubins HB, Robins SJ, Collins D, et al. Gemfibrozil for the secondary prevention of coronary heart disease in men with low levels of high-density lipoprotein cholesterol. Veterans Affairs High-Density Lipoprotein Cholesterol Intervention Trial Study Group. *N Engl J Med*. 1999;341:410–418. Kajioka T, Tsuzuku S, Shimokata H, Sato Y. Effects of intentional weight cycling on non-obese young women. *Metabolism*. 2002;51:149–154. Antic V, Tempini A, Montani JP. Serial changes in cardiovascular and renal function of rabbits ingesting a high-fat, high-calorie diet. *Am J Hypertens*. 1999;12:826–829. Schulz M, Liese AD, Boeing H, et al. Associations of short-term weight changes and weight cycling with incidence of essential hypertension in the EPIC-Potsdam Study. *J Hum Hypertens*. 2005;19:61–67.
4. Lagiou P, Sandin S, Lof M, et al. Low carbohydrate-high protein diet and incidence of cardiovascular diseases in Swedish women: prospective cohort study. *BMJ*. 2012;344:e4026. Lagiou P, Sandin S, Weiderpass E, et al. Low carbohydrate-high protein diet and mortality in a cohort of Swedish women. *J Intern Med*. 2007;261:366–374.
 5. Lagiou P, Sandin S, Lof M, et al: Low carbohydrate-high protein diet and incidence of cardiovascular diseases in Swedish women: prospective cohort study. *BMJ*. 2012;344:e4026. Lagiou P, Sandin S, Weiderpass E, et al: Low carbohydrate-high protein diet and mortality in a cohort of Swedish women. *J Intern Med*. 2007;261:366–374.
 6. Isner JM, Sours HE, Paris AL, et al. Sudden, unexpected death in avid dieters using the liquid-protein-modified-fast diet. Observations in 17 patients and the role of the prolonged QT interval. *Circulation*. 1979;60(6):1401–12.
 7. Stevens A, Robinson P, Turpin J et al. Sudden Cardiac Death of an Adolescent During [Atkins] *Dieting Southern Medical Journal*. 95(2002):1047.
 8. Tseng M, Olufade TO, Evers KA, Byrne C. Adolescent lifestyle factors and adult breast density in US Chinese immigrant women. *Nutr Cancer*. 2011; 63(3): 342–349. Berkey CS, Frazier AL, Gardner JD, Colditz GA. Adolescence and breast carcinoma risk. *Cancer*. 1999;85(11):2400–9.
 9. Cui X, Dai Q, Tseng M, et al. Dietary Patterns and Breast Cancer Risk in the Shanghai Breast Cancer Study. *Cancer Epidemiol Biomarkers Prev*. 2007;16:1443–1448.
 10. Reis JP, Loria CM, Lewis CE, et al. Association between duration of overall and abdominal obesity beginning in young adulthood and coronary artery calcification in middle age. *JAMA*. 2013;310(3):280–288.
 11. Calle EE, Kaaks R. Overweight, obesity and cancer: epidemiological evidence and proposed mechanisms. *Nat Rev Cancer*. 2004;4:579–591.
 12. Rastmanesh R. High polyphenol, low probiotic diet for weight loss because of intestinal microbiota interaction. *Chem Biol Interact*. 2011;189(1–2):1–8.

13. Jeffery IB, O'Toole PW. Diet-microbiota interactions and their implications for healthy living. *Nutrients*. 2013;5(1):234–252. Koeth RA, Wang Z, Levison BS, et al. Intestinal microbiota metabolism of l-carnitine, a nutrient in red meat, promotes atherosclerosis. *Nat Med*. 2013;19(5):576–585.
14. Andersen C, Rayalam S, Della-Fera MA, Baile CA. Phytochemicals and adipogenesis. *Biofactors*. 2010;36(6):415–422. Kim HK, Kim JN, Han SN, et al. Black soybean anthocyanins inhibit adipocyte differentiation in 3T3-L1 cells. *Nutr Res*. 2012;32(10):770–777. Rayalam S, Della-Fera MA, Baile CA. Phytochemicals and regulation of the adipocyte life cycle. *J Nutr Biochem*. 2008;19(11):717–726. Baile CA, Yang JY, Rayalam S, et al. Effect of resveratrol on fat mobilization. *Ann NY Acad Sci*. 2011;1215:40–47. Rayalam S, Della-Fera MA, Yang JY, et al. Resveratrol potentiates genistein's antiadipogenic and proapoptotic effects in 3T3-L1 adipocytes. *J Nutr*. 2007;137(12):2668–2673.

Четвърта глава: Силата на истинската храна

1. Moore SC, Carter LM, van Goozen S. Confectionery consumption in childhood and adult violence. *Br J Psychiatry*. 2009;195(4):366–367. Gesch CB. Influence of supplementary vitamins, minerals and essential fatty acids on the antisocial behaviour of young adult prisoners: Randomised, placebo-controlled trial. *The British Journal of Psychiatry*. 2002 2002/07/01;181(1):22–8. 7. Virkkunen M, Huttunen MO. Evidence for abnormal glucose tolerance test among violent offenders. *Neuropsychobiology*. 1982;8(1):30–4. PubMed PMID: 7057987. Golomb BA, Evans MA, White HL, et al. Trans fat consumption and aggression. *PloS one*. 2012;7(3):e32175. PubMed PMID: 22403632. Pubmed Central PMCID: 3293881. Yau PL, Castro MG, Tagani A, et al. Obesity and metabolic syndrome and functional and structural brain impairments in adolescence. *Pediatrics*. 2010;130:e856–864. Thaler JP, Yi C-X, Schur EA, et al. Obesity is associated with hypothalamic injury in rodents and humans. *Journal of Clinical Investigation*. 2012 2012/01/03;122(1):153–62. Northstone K, Joinson C, Emmett P, et al. Are dietary patterns in childhood associated with IQ at 8 years of age? A population-based cohort study. *Journal of Epidemiology and Community Health*. 2012 Jul;66(7):624–8. PubMed PMID: 21300993.
2. Science Daily. Consumers over age 50 should consider cutting copper and iron intake, report suggests. <http://www.sciencedaily.com/releases/2010/01/100120113553.htm>. Published January 22, 2010; accessed January 29, 2010. Brewer GJ. Risks of copper and iron toxicity during aging in humans. *Chem Res Toxicol*. 2010;23(2):319–326. Brewer GJ. Iron and copper toxicity in diseases of aging, particularly atherosclerosis and Alzheimer's disease. *Exp Biol Med*. 2007;232(2):323–335. Kidd PM. Neurodegeneration from mitochondrial insufficiency: Nutrients, stem cells, growth factors, and prospects for brain rebuilding using integrative management. *Alternative Medicine Review*. 2005;10(4):268–293. Morris MC, Evans DA, Tangney CC, et al. Dietary copper and high saturated

- and trans fat intakes associated with cognitive decline. *Arch Neurol*. 2006; 63(8):1085–1088.
3. Milton K. Back to basics: why foods of wild primates have relevance for modern human health. *Nutrition*. 2000;16(7/8):480–484.
 4. Richer SP. Randomized, double-blind, placebo-controlled study of zeaxanthin and visual function in patients with atrophic age-related macular degeneration. *Optometry*. 2011;82(11):667–680. Semba RD. Are lutein and zeaxanthin conditionally essential nutrients for eye health? *Med Hypotheses*. 2003;61(4):465–472. Willett W, Sampson L, Stampfer M, et al. Reproducibility and validity of a semiquantitative food frequency questionnaire. *Am J Epidemiol*. 1985;122:51–65. Delcourt C, Carrière I, Delage M, et al. Plasma lutein and zeaxanthin and other carotenoids as modifiable risk factors for age-related maculopathy and cataract: the POLA Study. *Invest Ophthalmol Vis Sci*. 2006;47:2329–2335.
 5. Higdon J, Delage B, Williams D, et al. Cruciferous vegetables and human cancer risk: epidemiologic evidence and mechanistic basis. *Pharmacol Res*. 2007;55:224–236.
 6. Cornblatt BS, Ye L, Dinkova-Kostova AT, et al. Preclinical and clinical evaluation of sulforaphane for chemoprevention in the breast. *Carcinogenesis*. 2007;28:1485–1490.
 7. Zhang CX, Ho SC, Chen YM, et al. Greater vegetable and fruit intake is associated with a lower risk of breast cancer among Chinese women. *Int J Cancer*. 2009;125:181–188.
 8. Bosetti C, Filomeno M, Riso P, et al. Cruciferous vegetables and cancer risk in a network of case-control studies. *Ann Oncol*. 2012;23(8):2198–2203.
 9. Nechuta SJ, Lu W, Cai H, et al. Cruciferous vegetable intake after diagnosis of breast cancer and survival: a report from the Shanghai Breast Cancer Survival study. Presented at the Annual Meeting of the American Association for Cancer Research; March 31–April 4, 2012; Chicago. Abstract no. LB-322.
 10. Thomson CA, Rock CL, Thompson PA, et al. Vegetable intake is associated with reduced breast cancer recurrence in tamoxifen users: a secondary analysis from the Women's Healthy Eating and Living Study. *Breast Cancer Res Treat*. 2011;125:519–527.
 11. Garg M, Garg C, Mukherjee PK, Suresh B. Antioxidant potential of *Lactuca sativa*. *Anc Sci Life*. 2004;24(1):6–10. Mulabagala V, Ngouajob M, Nairb A, et al. In vitro evaluation of red and green lettuce (*Lactuca sativa*) for functional food properties. *Food Chem*. 2010;118(2):300–306. Serafini M, Bugianesi R, Salucci M, et al. Effect of acute ingestion of fresh and stored lettuce (*Lactuca sativa*) on plasma total antioxidant capacity and antioxidant levels in human subjects. *Br J Nutr*. 2002;88(6):615–623. Gridling M, Popescu R, Kopp B, et al. Anti-leukaemic effects of two extract types of

- Lactuca sativa correlate with the activation of Chk2, induction of p21, downregulation of cyclin D1 and acetylation of alpha-tubulin. *Oncol Rep.* 2010;23(4):1145–1151.
12. Zhao J, Moore AN, Redell JB, et al. Enhancing expression of Nrf2-driven genes protects the blood brain barrier after brain injury. *J Neurosci.* 2007;27:10240–10248. Carter P, Gray LJ, Troughton J, et al. Fruit and vegetable intake and incidence of type 2 diabetes mellitus: systematic review and meta-analysis. *BMJ.* 2010;341:c4229. Lundberg JO, Carlstrom M, Larsen FJ, et al. Roles of dietary inorganic nitrate in cardiovascular health and disease. *Cardiovasc Res.* 2011;89:525–532. Higdon J, Delage B, Williams D, et al. Cruciferous vegetables and human cancer risk: epidemiologic evidence and mechanistic basis. *Pharmacol Res.* 2007;55:224–236. Stringham JM, Bovier ER, Wong JC, et al. The influence of dietary lutein and zeaxanthin on visual performance. *J Food Sci.* 2010;75:R24–R29.
 13. Mulabagala V, Ngouajjob M, Nairb A, et al. In vitro evaluation of red and green lettuce (*Lactuca sativa*) for functional food properties. *Food Chem.* 2010;118(2):300–306.
 14. World Cancer Research Fund, American Institute for Cancer Research. Expert report. Food, nutrition, physical activity, and the prevention of cancer: a global perspective. World Cancer Research Fund; 2007.
 15. Bessaoud F, Daurès JP, Gerber M. Dietary factors and breast cancer risk: a case control study among a population in Southern France. *Nutr Cancer.* 2008;60(2):177–187. Link LB, Potter JD. Raw versus cooked vegetables and cancer risk. *Cancer Epidemiol Biomarkers Prev.* 2004;13(9):1422–1435.
 16. Cox BD, Whichelow MJ, Prevost AT. Seasonal consumption of salad vegetables and fresh fruit in relation to the development of cardiovascular disease and cancer. *Public Health Nutr.* 2000;3(1):19–29. Lockheart MS, Steffen LM, Rebnord HM, et al. Dietary patterns, food groups and myocardial infarction: a case-control study. *Br J Nutr.* 2007;98(2):380–387. Oude Griep LM, Verschuren WM, Kromhout D, et al. Raw and processed fruit and vegetable consumption and 10-year stroke incidence in a population-based cohort study in the Netherlands. *Eur J Clin Nutr.* 2011;65(7):791–799.
 17. Rolls BJ, Roe LS, Meengs JS. Salad and satiety: energy density and portion size of a first-course salad affect energy intake at lunch. *J Am Diet Assoc.* 2004;104(10):1570–1576. Roe LS, Meengs JS, Rolls BJ. Salad and satiety: the effect of timing of salad consumption on meal energy intake. *Appetite.* 2012;58(1):242–248.
 18. Behall KM, Howe JC. Effect of long-term consumption of amylose vs amylopectin starch on metabolic variables in human subjects. *Am J Clin Nutr.* 1995;61:334–340. Marlett JA, McBurney MI, Slavin JL. Position of the American Dietetic Association: health implications of dietary fiber. *J Am Diet Assoc.* 2002;102:993–1000. Aller EE, Abete I, Astrup A, et al. Starches, sugars and obesity. *Nutrients.* 2011;3:341–369. Wylie-Rosett J, Segal-Isaacson CJ,

- Segal-Isaacson A. Carbohydrates and increases in obesity: does the type of carbohydrate make a difference? *Obes Res.* 2004;12(suppl 2):124S–129S.
19. Bednar GE, Patil AR, Murray SM, et al. Starch and fiber fractions in selected food and feed ingredients affect their small intestinal digestibility and fermentability and their large bowel fermentability in vitro in a canine model. *J Nutr.* 2001;131:276–286. Muir JG, O’Dea K. Measurement of resistant starch: factors affecting the amount of starch escaping digestion in vitro. *Am J Clin Nutr.* 1992;56:123–127.
 20. Jenkins DJ, Kendall CW, Augustin LS, et al. Effect of legumes as part of a low glycemic index diet on glycemic control and cardiovascular risk factors in type 2 diabetes mellitus: a randomized controlled trial. *Arch Intern Med.* 2012;172(21):1653–1660.
 21. Halton TL, Willett WC, Liu S, et al. Potato and french fry consumption and risk of type 2 diabetes in women. *Am J Clin Nutr.* 2006;83:284–290.
 22. Streppel MT, Arends LR, van’t Veer P, et al. Dietary fiber and blood pressure: a meta-analysis of randomized placebo-controlled trials. *Arch Intern Med.* 2005;165:150–156. Houston MC. The importance of potassium in managing hypertension. *Curr Hypertens Rep.* 2011;13:309–317. Houston M. The role of magnesium in hypertension and cardiovascular disease. *J Clin Hypertens.* 2011;13:843–847. DeFronzo RA, Cooke CR, Andres R, et al. The effect of insulin on renal handling of sodium, potassium, calcium, and phosphate in man. *J Clin Invest.* 1975;55:845–855. Chiasson JL, Josse RG, Gomis R, et al. Acarbose treatment and the risk of cardiovascular disease and hypertension in patients with impaired glucose tolerance: the STOP-NIDDM trial. *JAMA.* 2003;290:486–494.
 23. Papanikolaou Y, Fulgoni VL. Bean consumption is associated with greater nutrient intake, reduced systolic blood pressure, lower body weight, and a smaller waist circumference in adults: results from the National Health and Nutrition Examination Survey 1999–2002. *J Am Coll Nutr.* 2008;27:569–576. Bazzano LA, Thompson AM, Tees MT, et al. Non-soy legume consumption lowers cholesterol levels: a meta-analysis of randomized controlled trials. *Nutr Metab Cardiovasc Dis.* 2011;21:94–103.
 24. Hamer HM, Jonkers D, Venema K, et al. Review article: the role of butyrate on colonic function. *Aliment Pharmacol Ther.* 2008;27:104–119. O’Keefe SJ, Ou J, Aufreiter S, et al. Products of the colonic microbiota mediate the effects of diet on colon cancer risk. *J Nutr.* 2009;139:2044–2048. Singh PN, Fraser GE. Dietary risk factors for colon cancer in a low-risk population. *Am J Epidemiol.* 1998;148:761–774.
 25. Aune D, De Stefani E, Ronco A, et al. Legume intake and the risk of cancer: a multisite case-control study in Uruguay. *Cancer Causes Control.* 2009;20:1605–1615. Bao PP, Shu XO, Zheng Y, et al. Fruit, vegetable, and animal food intake and breast cancer risk by hormone receptor status. *Nutr Cancer.* 2012;64:806–819.

26. Dong JY, He K, Wang P, Qin LQ. Dietary fiber intake and risk of breast cancer: a meta-analysis of prospective cohort studies. *Am J Clin Nutr.* 2011;94(3):900–905.
27. Powolny A, Singh S. Multitargeted prevention and therapy of cancer by diallyl trisulfide and related Allium vegetable-derived organosulfur compounds. *Cancer Lett.* 2008;269:305–314. Ginter E, Simko V. Garlic (*Allium sativum* L.) and cardiovascular diseases. *Bratisl Lek Listy.* 2010;111:452–456. Taj Eldin IM, Ahmed EM, Elwahab HMA. Preliminary study of the clinical hypoglycemic effects of *Allium cepa* (red onion) in type 1 and type 2 diabetic patients. *Environ Health Insights.* 2010;4:71–77.
28. Slimestad R, Fossen T, Vagen IM. Onions: a source of unique dietary flavonoids. *J Agric Food Chem.* 2007;55:10067–10080. Miyamoto S, Yasui Y, Ohigashi H, et al. Dietary flavonoids suppress azoxymethane-induced colonic preneoplastic lesions in male C57BL/KsJ-db/db mice. *Chem Biol Interact.* 2010;183:276–283. Shan BE, Wang MX, Li RQ. Quercetin inhibits human SW480 colon cancer growth in association with inhibition of cyclin D1 and survivin expression through Wnt/beta-catenin signaling pathway. *Cancer Invest.* 2009;27:604–612.
29. Galeone C, Pelucchi C, Levi F, et al. Onion and garlic use and human cancer. *Am J Clin Nutr.* 2006;84:1027–1032.
30. Modem S, Dicarolo SE, Reddy TR. Fresh garlic extract induces growth arrest and morphological differentiation of MCF7 breast cancer cells. *Genes Cancer.* 2012;3:177–186. Na HK, Kim EH, Choi MA, et al. Diallyl trisulfide induces apoptosis in human breast cancer cells through ROS-mediated activation of JNK and AP-1. *Biochem Pharmacol.* 2012;84(10):1241–1250. Malki A, El-Saadani M, Sultan AS. Garlic constituent diallyl trisulfide induced apoptosis in MCF7 human breast cancer cells. *Cancer Biol Ther.* 2009;8:2175–2185.
31. Ravasco P, Aranha MM, Borralho PM, et al. Colorectal cancer: can nutrients modulate NF-kappaB and apoptosis? *Clin Nutr.* 2010;29:42–46. Pierini R, Gee JM, Belshaw NJ, et al. Flavonoids and intestinal cancers. *Br J Nutr.* 2008;99(suppl 1):ES53–59.
32. Yu L, Fernig DG, Smith JA, et al. Reversible inhibition of proliferation of epithelial cell lines by *Agaricus bisporus* (edible mushroom) lectin. *Cancer Res* 1993;53:4627–4632.
33. Borchers AT, Krishnamurthy A, Keen CL, et al. The immunobiology of mushrooms. *Exp Biol Med.* 2008;233:259–276.
34. Zhang M, Huang J, Xie X, et al. Dietary intakes of mushrooms and green tea combine to reduce the risk of breast cancer in Chinese women. *Int J Cancer.* 2009;124:1404–1408.
35. Hong SA, Kim K, Nam SJ, et al. A case-control study on the dietary intake of mushrooms and breast cancer risk among Korean women. *Int J Cancer.* 2008;122:919–923. Shin A, Kim J, Lim SY, et al. Dietary mushroom intake

- and the risk of breast cancer based on hormone receptor status. *Nutr Cancer*. 2010;62:476–483.
36. Cao QZ, Lin ZB. Antitumor and anti-angiogenic activity of *Ganoderma lucidum* polysaccharides peptide. *Acta Pharmacol Sin*. 2004;25:833–838. Lee JS, Park BC, Ko YJ, et al. *Grifola frondosa* (maitake mushroom) water extract inhibits vascular endothelial growth factor-induced angiogenesis through inhibition of reactive oxygen species and extracellular signal-regulated kinase phosphorylation. *J Med Food*. 2008;11:643–651. Chang HH, Hsieh KY, Yeh CH, et al. Oral administration of an Enoki mushroom protein FVE activates innate and adaptive immunity and induces anti-tumor activity against murine hepatocellular carcinoma. *Int Immunopharmacol*. 2010;10:239–246.
 37. Hara M, Hanaoka T, Kobayashi M, et al. Cruciferous vegetables, mushrooms, and gastrointestinal cancer risks in a multicenter, hospital-based case-control study in Japan. *Nutr Cancer*. 2003;46:138–147. Zhang CX, Ho SC, Chen YM, et al. Greater vegetable and fruit intake is associated with a lower risk of breast cancer among Chinese women. *Int J Cancer*. 2009;125:181–188. Martin KR, Brophy SK. Commonly consumed and specialty dietary mushrooms reduce cellular proliferation in MCF-7 human breast cancer cells. *Exp Biol Med*. 2010;235:1306–1314. Fang N, Li Q, Yu S, et al. Inhibition of growth and induction of apoptosis in human cancer cell lines by an ethyl acetate fraction from shiitake mushrooms. *J Altern Complement Med*. 2006;12:125–132. Ng ML, Yap AT. Inhibition of human colon carcinoma development by lentinan from shiitake mushrooms (*Lentinus edodes*). *J Altern Complement Med*. 2002;8:581–589. Adams LS, Phung S, Wu X, et al. White button mushroom (*Agaricus bisporus*) exhibits antiproliferative and proapoptotic properties and inhibits prostate tumor growth in athymic mice. *Nutr Cancer*. 2008;60:744–756. Lakshmi B, Ajith TA, Sheena N, et al. Antiperoxidative, anti-inflammatory, and antimutagenic activities of ethanol extract of the mycelium of *Ganoderma lucidum* occurring in South India. *Teratog Carcinog Mutagen*. 2003;suppl 1:85–97. Lin ZB, Zhang HN. Anti-tumor and immunoregulatory activities of *Ganoderma lucidum* and its possible mechanisms. *Acta Pharmacol Sin*. 2004;25:1387–1395.
 38. Poddar KH, Ames M, Chen H, et al. Positive effect of white button mushrooms when substituted for meat on body weight and composition changes during weight loss and weight maintenance: a one-year randomized clinical trial. *FASEB J*. 2013;27:852.4.
 39. Su CH, Lai MN, Ng LT. Inhibitory effects of medicinal mushrooms on α -amylase and α -glucosidase enzymes related to hyperglycemia. *Food Funct*. 2013;4(4):644–649.
 40. Varshney J, Ooi JH, Jayarao BM, et al. White button mushrooms increase microbial diversity and accelerate the resolution of *Citrobacter rodentium* infection in mice. *J Nutr*. 2013;143(4):526–532.

41. Jeong SC, Koyyalamudi SR, Pang G. Dietary intake of *Agaricus bisporus* white button mushroom accelerates salivary immunoglobulin A secretion in healthy volunteers. *Nutrition*. 2012;28(5):527–531.
42. Toth B, Erickson J. Cancer induction in mice by feeding of the uncooked cultivated mushroom of commerce *Agaricus bisporus*. *Cancer Res*. 1986;46:4007–4011. Schulzova V, Hajslova J, Peroutka R, et al. Influence of storage and household processing on the agaritine content of the cultivated *Agaricus* mushroom. *Food Addit Contam*. 2002;19:853–862.
43. Huxley RR, Neil HA. The relation between dietary flavonol intake and coronary heart disease mortality: a meta-analysis of prospective cohort studies. *Eur J Clin Nutr*. 2003;57:904–908. Knekt P, Kumpulainen J, Jarvinen R, et al. Flavonoid intake and risk of chronic diseases. *Am J Clin Nutr*. 2002;76:560–568. Mursu J, Voutilainen S, Nurmi T, et al. Flavonoid intake and the risk of ischaemic stroke and CVD mortality in middle-aged Finnish men: the Kuopio Ischaemic Heart Disease Risk Factor Study. *Brit J Nutr*. 2008;100:890–895. Mink PJ, Scrafford CG, Barraj LM, et al. Flavonoid intake and cardiovascular disease mortality: a prospective study in postmenopausal women. *Am J Clin Nutr*. 2007;85:895–909.
44. Bazzano LA, Li TY, Joshipura KJ, et al. Intake of fruit, vegetables, and fruit juices and risk of diabetes in women. *Diabetes Care*. 2008;31:1311–1317. Hannum SM. Potential impact of strawberries on human health: a review of the science. *Crit Rev Food Sci Nutr*. 2004;44:1–17. Joseph JA, Shukitt-Hale B, Willis LM. Grape juice, berries, and walnuts affect brain aging and behavior. *J Nutr*. 2009;139:1813S–1817S. Cassidy A, O'Reilly EJ, Kay C, et al. Habitual intake of flavonoid subclasses and incident hypertension in adults. *Am J Clin Nutr*. 2011;93:338–347. Devore EE, Kang JH, Breteler MM, et al. Dietary intakes of berries and flavonoids in relation to cognitive decline. *Ann Neurol*. 2012;72(1):135–143.
45. Smart RC, Huang MT, Chang RL, et al. Disposition of the naturally occurring antimutagenic plant phenol, ellagic acid, and its synthetic derivatives, 3-O-decylellagic acid and 3,3'-di-O-methylellagic acid in mice. *Carcinogenesis*. 1986;7:1663–1667. Smart RC, Huang MT, Chang RL, et al. Effect of ellagic acid and 3-O-decylellagic acid on the formation of benzo[a]pyrene-derived DNA adducts in vivo and on the tumorigenicity of 3-methylcholanthrene in mice. *Carcinogenesis*. 1986;7:1669–1675.
46. Stoner GD, Wang LS, Casto BC. Laboratory and clinical studies of cancer chemoprevention by antioxidants in berries. *Carcinogenesis*. 2008;29:1665–1674.
47. Kim ND, Mehta R, Yu W, et al. Chemopreventive and adjuvant therapeutic potential of pomegranate (*Punica granatum*) for human breast cancer. *Breast Cancer Res Treat*. 2002;71:203–217. Kohno H, Suzuki R, Yasui Y, et al. Pomegranate seed oil rich in conjugated linolenic acid suppresses chemically induced colon carcinogenesis in rats. *Cancer Sci*. 2004;95:481–486.

- Kawaii S, Lansky EP. Differentiation-promoting activity of pomegranate (*Punica granatum*) fruit extracts in HL-60 human promyelocytic leukemia cells. *J Med Food*. 2004;7:13–18.
48. Roy S, Khanna S, Alessio HM, et al. Anti-angiogenic property of edible berries. *Free Radic Res*. 2002;36:1023–1031. Khan N, Afaq F, Kweon MH, et al. Oral consumption of pomegranate fruit extract inhibits growth and progression of primary lung tumors in mice. *Cancer Res*. 2007;67:3475–3482. Toi M, Bando H, Ramachandran C, et al. Preliminary studies on the anti-angiogenic potential of pomegranate fractions in vitro and in vivo. *Angiogenesis*. 2003;6:121–128. Sartippour MR, Seeram NP, Rao JY, et al. Ellagitannin-rich pomegranate extract inhibits angiogenesis in prostate cancer in vitro and in vivo. *Int J Oncol*. 2008;32:475–480. Panchal SK, Ward L, Brown L. Ellagic acid attenuates high-carbohydrate, high-fat diet-induced metabolic syndrome in rats. *Eur J Nutr*. 2013;52(2):559–568. Edirisinghe I, Banaszewski K, Capozzo J, et al. Strawberry anthocyanin and its association with postprandial inflammation and insulin. *Br J Nutr*. 2011;106:913–922. Adams LS, Seeram NP, Aggarwal BB, et al. Pomegranate juice, total pomegranate ellagitannins, and punicalagin suppress inflammatory cell signaling in colon cancer cells. *J Agric Food Chem*. 2006;54:980–985.
 49. Adams LS, Zhang Y, Seeram NP, et al. Pomegranate ellagitannin-derived compounds exhibit antiproliferative and antiaromatase activity in breast cancer cells in vitro. *Cancer Prev Res*. 2010;3:108–113.
 50. American Association for Cancer Research. Strawberries may slow precancerous growth in esophagus. <http://aacrnews.wordpress.com/2011/04/06/strawberries-may-slow-precancerous-growth-in-esophagus/>. Published April 6, 2011.
 51. Boqué N, de la Iglesia R, de la Garza AL, et al. Prevention of diet-induced obesity by apple polyphenols in Wistar rats through regulation of adipocyte gene expression and DNA methylation patterns. *Mol Nutr Food Res*. 2013;57(8):1473–1478. doi: 10.1002/mnfr.201200686. Chang JJ, Hsu MJ, Huang HP, et al. Mulberry anthocyanins inhibit oleic acid induced lipid accumulation by reduction of lipogenesis and promotion of hepatic lipid clearance. *J Agric Food Chem*. 2013;61(25):6069–6076.
 52. Jeffery IB, O'Toole PW. Diet-microbiota interactions and their implications for healthy living. *Nutrients*. 2013;5(1):234–52. Tuohy KM, Conterno L, Gasperotti M, Viola R. Up-regulating the human intestinal microbiome using whole plant foods, polyphenols, and/or fiber. *J Agric Food Chem*. 2012;60(36):8776–82. Pray L, Pillsbury L, Tomayko E, The Human Microbiome, Diet, and Health: Workshop Summary Institute of Medicine (US) Food Forum. Food and Nutrition Board (FNB) Washington (DC): National Academies Press (US); 2013.
 53. Cassidy A, O'Reilly EJ, Kay C, et al. Habitual intake of flavonoid subclasses and incident hypertension in adults. *Am J Clin Nutr*. 2011;93(2):338–347.

54. Cassidy A, Mukamal KJ, Liu L, et al. High anthocyanin intake is associated with a reduced risk of myocardial infarction in young and middle-aged women. *Circulation*. 2013;127:188–196.
55. Galleano M, Pechanova O, Fraga CG. Hypertension, nitric oxide, oxidants, and dietary plant polyphenols. *Curr Pharm Biotechnol*. 2010;11:837–848. Basu A, Rhone M, Lyons TJ. Berries: emerging impact on cardiovascular health. *Nutr Rev*. 2010;68:168–177. Chong MF, Macdonald R, Lovegrove JA. Fruit polyphenols and CVD risk: a review of human intervention studies. *Brit J Nutr*. 2010;104(suppl 3):S28–S39. Basu A, Du M, Leyva MJ, et al. Blueberries decrease cardiovascular risk factors in obese men and women with metabolic syndrome. *J Nutr*. 2010;140:1582–1587.
56. Devore EE, Kang JH, Breteler MM, et al. Dietary intakes of berries and flavonoids in relation to cognitive decline. *Ann Neurol*. 2012;72(1):135–143.
57. Joseph JA, Shukitt-Hale B, Willis LM. Grape juice, berries, and walnuts affect brain aging and behavior. *J Nutr*. 2009;139:1813S–1817S. Ma Y, Njike VY, Millet J, et al. Effects of walnut consumption on endothelial function in type 2 diabetic subjects: a randomized controlled crossover trial. *Diabetes Care*. 2010;33:227–232.
58. Mattes RD, Dreher ML. Nuts and healthy body weight maintenance mechanisms. *Asia Pac J Clin Nutr*. 2010;19:137–141.
59. Thompson LU, Chen JM, Li T, et al. Dietary flaxseed alters tumor biological markers in postmenopausal breast cancer. *Clin Cancer Res*. 2005;11:3828–3835.
60. McCann SE, Thompson LU, Nie J, et al. Dietary lignan intakes in relation to survival among women with breast cancer: the Western New York Exposures and Breast Cancer (WEB) Study. *Breast Cancer Res Treat*. 2010;122(1):229–235.
61. Connor WE. Importance of n-3 fatty acids in health and disease. *Am J Clin Nutr*. 2000;71(suppl 1):171S–175S.
62. Yehuda S, Rabinovitz S, Mostofsky DI. Mixture of essential fatty acids lowers test anxiety. *Nutr Neurosci*. 2005;8(4):265–267. Lucas M, Mirzaei F, O'Reilly EJ, et al. Dietary intake of n-3 and n-6 fatty acids and the risk of clinical depression in women: a 10-y prospective follow-up study. *Am J Clin Nutr*. 2011;93(6):1337–1343.
63. Shardell MD, Alley DE, Hicks GE, et al. Low-serum carotenoid concentrations and carotenoid interactions predict mortality in US adults: the Third National Health and Nutrition Examination Survey. *Nutr Res*. 2011;31:178–189.
64. Canene-Adams K, Campbell JK, Zaripheh S, et al. The tomato as a functional food. *J Nutr*. 2005;135:1226–1230.

65. van Breemen RB, Pajkovic N. Multitargeted therapy of cancer by lycopene. *Cancer Lett.* 2008;269:339–351.
66. Rizwan M, Rodriquez-Blanco I, Harbottle A, et al. Tomato paste rich in lycopene protects against cutaneous photodamage in humans in vivo. *Br J Dermatol.* 2011;164(1):154–162.
67. Rissanen TH, Voutilainen S, Nyyssonen K, et al. Low serum lycopene concentration is associated with an excess incidence of acute coronary events and stroke: the Kuopio Ischaemic Heart Disease Risk Factor Study. *Br J Nutr.* 2001;85:749–754. Rissanen T, Voutilainen S, Nyyssonen K, et al. Lycopene, atherosclerosis, and coronary heart disease. *Exp Biol Med.* 2002;227:900–907. Rissanen TH, Voutilainen S, Nyyssonen K, et al. Serum lycopene concentrations and carotid atherosclerosis: the Kuopio Ischaemic Heart Disease Risk Factor Study. *Am J Clin Nutr.* 2003;77:133–138.
68. Sesso HD, Buring JE, Norkus EP, et al. Plasma lycopene, other carotenoids, and retinol and the risk of cardiovascular disease in women. *Am J Clin Nutr.* 2004;79:47–53.
69. Hak AE, Ma J, Powell CB, et al. Prospective study of plasma carotenoids and tocopherols in relation to risk of ischemic stroke. *Stroke.* 2004;35:1584–1588.
70. Karppi J, Laukkanen JA, Sivenius J, et al. Serum lycopene decreases the risk of stroke in men: a population-based follow-up study. *Neurology.* 2012;79:1540–1547.
71. Karppi J, Laukkanen JA, Makikallio TH, et al. Low serum lycopene and beta-carotene increase risk of acute myocardial infarction in men. *Eur J Public Health.* 2012;22(6):835–840.
72. Silaste ML, Alfthan G, Aro A, et al. Tomato juice decreases LDL cholesterol levels and increases LDL resistance to oxidation. *Br J Nutr.* 2007;98:1251–1258. Burton-Freeman B, Talbot J, Park E, et al. Protective activity of processed tomato products on postprandial oxidation and inflammation: a clinical trial in healthy weight men and women. *Mol Nutr Food Res.* 2012;56:622–631. Hadley CW, Clinton SK, Schwartz SJ. The consumption of processed tomato products enhances plasma lycopene concentrations in association with a reduced lipoprotein sensitivity to oxidative damage. *J Nutr.* 2003;133:727–732.
73. Xaplanteris P, Vlachopoulos C, Pietri P, et al. Tomato paste supplementation improves endothelial dynamics and reduces plasma total oxidative status in healthy subjects. *Nutr Res.* 2012;32:390–394.
74. Ried K, Fakler P. Protective effect of lycopene on serum cholesterol and blood pressure: meta-analyses of intervention trials. *Maturitas.* 2011;68:299–310.

75. Palozza P, Parrone N, Catalano A, et al. Tomato lycopene and inflammatory cascade: basic interactions and clinical implications. *Curr Med Chem.* 2010;17:2547–2563. Palozza P, Parrone N, Simone RE, et al. Lycopene in atherosclerosis prevention: an integrated scheme of the potential mechanisms of action from cell culture studies. *Arch Biochem Biophys.* 2010;504:26–33.
76. USDA, Economic Research Service. Food availability (per capita) data system. [http://www.ers.usda.gov/data-products/food-availability-\(per-capita\)-data-system.aspx](http://www.ers.usda.gov/data-products/food-availability-(per-capita)-data-system.aspx).
77. Brown MJ, Ferruzzi MG, Nguyen ML, et al. Carotenoid bioavailability is higher from salads ingested with full-fat than with fat-reduced salad dressings as measured with electrochemical detection. *Am J Clin Nutr.* 2004;80:396–403.
78. van het Hof KH, de Boer BC, Tijburg LB, et al. Carotenoid bioavailability in humans from tomatoes processed in different ways determined from the carotenoid response in the triglyceride-rich lipoprotein fraction of plasma after a single consumption and in plasma after four days of consumption. *J Nutr.* 2000;130:1189–1196. Goltz SR, Campbell WW, Chitchumroonchokchai C, et al. Meal triacylglycerol profile modulates postprandial absorption of carotenoids in humans. *Mol Nutr Food Res.* 2012;56:866–877.

Пета глава: Начално обучение за нутритарианци

1. USDA, Economic Research Service. Food availability (per capita) data system. [http://www.ers.usda.gov/data-products/food-availability-\(per-capita\)-data-system.aspx#.Ud8A6UG1Gck](http://www.ers.usda.gov/data-products/food-availability-(per-capita)-data-system.aspx#.Ud8A6UG1Gck). Lin B-H, Yen ST. The U.S. grain consumption landscape: who eats grain, in what form, where, and how much. USDA Economic Research Service; Economic Research Report No. 50; November 2007; http://www.ers.usda.gov/media/216648/err50_1_.pdf.
2. Halton TL, Willett WC, Liu S, et al. Potato and french fry consumption and risk of type 2 diabetes in women. *Am J Clin Nutr.* 2006;83(2):284–290.
3. Gnagnarella P, Gandini S, La Vecchia C, Maisonneuve P. Glycemic index, glycemic load, and cancer risk: a meta-analysis. *Am J Clin Nutr.* 2008;87(6):1793–1801. Barclay AW, Petocz P, McMillan-Price J, et al. Glycemic index, glycemic load, and chronic disease risk—a meta-analysis of observational studies. *Am J Clin Nutr.* 2008;87(3):627–637.
4. Williams CD, Satia JA, Adair LS, et al. Dietary patterns, food groups, and rectal cancer risk in Whites and African-Americans. *Cancer Epidemiol Biomarkers Prev.* 2009;18(5):1552–1561.
5. Sieri S, Krogh V, Berrino F, et al. Dietary glycemic load and index and risk of coronary heart disease in a large Italian cohort: the EPICOR study. *Arch Intern Med.* 2010;170(7):640–647.

6. Note TK.
7. Holt S, Brand Miller J, Petocz P. An insulin index of foods: the insulin demand generated by 1000-Kj portions of common foods. *Am J Clin Nutr.* 1997;66:1264–1276. Bao J, de Jong V, Atkinson F, et al. Food insulin index: physiologic basis for predicting insulin demand evoked by composite meals. *Am J Clin Nutr.* 2009;90:986–992.
8. Vergnaud AC, Norat T, Romaguera D, et al. Meat consumption and prospective weight change in participants of the EPIC-PANACEA study 1,2,3. *Am J Clin Nutr.* 2010;92(2):398–407. Sabaté J, Ang Y. Nuts and health outcomes: new epidemiologic evidence. *Am J Clin Nutr.* 2009;89(5):1643S–1648S. Fraser GE, Sabate J, Beeson WL, Strahan TM. A possible protective effect of nut consumption on risk of coronary heart disease: the Adventist Health Study. *Arch Intern Med.* 1992;152(7):1416–1424. Griel AE, Kris-Etherton PM. Tree nuts and the lipid profile: a review of clinical studies. *Br J Nutr.* 2006;96(suppl 2):S68–78. Sabaté J, Oda K, Ros E. Nut consumption and blood lipid levels: a pooled analysis of 25 intervention trials. *Arch Intern Med.* 2010;170(9):821–827. Mattes RD, Kris-Etherton PM, Foster GD. Impact of peanuts and tree nuts on body weight and healthy weight loss in adults. *J Nutr.* 2008;138:1741S–1745S. Bes-Rastrollo M, Sabate J, Gomez-Gracia E, et al. Nut consumption and weight gain in a Mediterranean cohort: the SUN study. *Obesity.* 2007;15:107–116. Bes-Rastrollo M, Wedick NM, Martinez-Gonzalez MA, et al. Prospective study of nut consumption, long-term weight change, and obesity risk in women. *Am J Clin Nutr.* 2009;89:1913–1919.
9. Chitnis MM, Yuen JS, Protheroe AS, et al. The type 1 insulin-like growth factor receptor pathway. *Clin Cancer Res.* 2008;14:6364–6370. Werner H, Bruchim I. The insulin-like growth factor-I receptor as an oncogene. *Arch Physiol Biochem.* 2009;115:58–71. Davies M, Gupta S, Goldspink G, et al. The insulin-like growth factor system and colorectal cancer: clinical and experimental evidence. *Int J Colorectal Dis.* 2006;21:201–208. Sandhu MS, Dunger DB, Giovannucci EL. Insulin, insulin-like growth factor-I (IGF-I), IGF binding proteins, their biologic interactions, and colorectal cancer. *J Natl Cancer Inst.* 2002;94:972–980. Rowlands MA, Gunnell D, Harris R, Vatten LJ, Holly JM, Martin RM. Circulating insulin-like growth factor peptides and prostate cancer risk: a systematic review and meta-analysis. *Int J Cancer.* 2009;124(10):2416–2429.
10. Werner H, Bruchim I. The insulin-like growth factor-1 receptor as an oncogene. *Arch Physiol Biochem.* 2009;115(2):58–71. Chitnis MM, Yuen JS, Protheroe AS, et al. The type 1 insulin-like growth factor receptor pathway. *Clin Cancer Res.* 2008;14(20):6364–6370.
11. Bartke A. Minireview: role of the growth hormone/insulin-like growth factor system in mammalian aging. *Endocrinology.* 2005;146:3718–3723.

12. Rinaldi S, Peeters PH, Berrino F, et al. IGF-I, IGFBP-3 and breast cancer risk in women: the European Prospective Investigation into Cancer and Nutrition (EPIC). *Endocr Relat Cancer*. 2006;13:593–605.
13. Hankinson SE, Willett WC, Colditz GA, et al. Circulating concentrations of insulin-like growth factor-I and risk of breast cancer. *Lancet*. 1998;351:1393–1396.
14. Lann D, LeRoith D. The role of endocrine insulin-like growth factor-I and insulin in breast cancer. *J Mammary Gland Biol Neoplasia*. 2008;13:371–379. Allen NE, Roddam AW, Allen DS, et al. A prospective study of serum insulin-like growth factor-I (IGF-I), IGF-II, IGF-binding protein-3 and breast cancer risk. *Br J Cancer*. 2005;92:1283–1287. Fletcher O, Gibson L, Johnson N, et al. Polymorphisms and circulating levels in the insulin-like growth factor system and risk of breast cancer: a systematic review. *Cancer Epidemiol Biomarkers Prev*. 2005;14:2–19. Renehan AG, Zwahlen M, Minder C, et al. Insulin-like growth factor (IGF)-I, IGF binding protein-3, and cancer risk: systematic review and meta-regression analysis *Lancet*. 2004;363(9418):1346–1353. Shi R, Yu H, McLarty J, et al. IGF-I and breast cancer: a meta-analysis. *Int J Cancer*. 2004;111(3):418–423. Sugumar A, Liu YC, Xia Q, et al. Insulin-like growth factor (IGF)-I and IGF-binding protein 3 and the risk of premenopausal breast cancer: a meta-analysis of literature. *Int J Cancer*. 2004;111(2):293–297. Key TJ, Appleby PN, Reeves GK, Roddam AW, Endogenous Hormones and Breast Cancer Collaborative Group. Insulin-like growth factor 1 (IGF1), IGF binding protein 3 (IGFBP3), and breast cancer risk: pooled individual data analysis of 17 prospective studies. *Lancet Oncol*. 2010;11(6):530–542.
15. Vardy ER, Rice PJ, Bowie PC, et al. Increased circulating insulin-like growth factor-1 in late-onset Alzheimer's disease. *J Alzheimers Dis*. 2007;12(4):285–290. Cohen E. Countering neurodegeneration by reducing the activity of the insulin/IGF signaling pathway: current knowledge and future prospects. *Exp Gerontol*. 2011;46(2–3):124–128.
16. Thissen JP, Ketelslegers JM, Underwood LE. Nutritional regulation of the insulin-like growth factors. *Endocr Rev*. 1994;15:80–101. Tsilidis KK, Travis RC, Appleby PN, et al. Insulin-like growth factor pathway genes and blood concentrations, dietary protein and risk of prostate cancer in the NCI Breast and Prostate Cancer Cohort Consortium (BPC3). *Int J Cancer*. 2013;133(2):495–504.
17. Qun LQ, He K, Xu JY. Milk consumption and circulating insulin-like growth factor-1 levels: a systematic literature review. *Int J Food Sci Nutr*. 2009;60(suppl 7):330–340. Heaney RP, McCarron DA, Dawson-Hughes B, et al. Dietary changes favorably affect bone remodeling in older adults. *J Am Diet Assoc*. 1999;99:1228–1233.
18. Roddam AW, Allen NE, Appleby P, et al. Insulin-like growth factors, their binding proteins, and prostate cancer risk: analysis of individual patient

- data from 12 prospective studies. *Ann Intern Med.* 2008;149(7):461–471. Sim HG, Cheng CW. Changing demography of prostate cancer in Asia. *Eur J Cancer.* 2005;41(6):834–845.
19. Song Y, Chavarro JE, Cao Y, et al. Whole milk intake is associated with prostate cancer-specific mortality among U.S. male physicians *J Nutr.* 2013;143(2):189–196.
 20. Giovannucci E, Pollak M, Liu Y, et al. Nutritional predictors of insulin-like growth factor I and their relationships to cancer in men. *Cancer Epidemiol Biomarkers Prev.* 2003;12:84–89. Holmes MD, Pollak MN, Willett WC, et al. Dietary correlates of plasma insulin-like growth factor I and insulin-like growth factor binding protein 3 concentrations. *Cancer Epidemiol Biomarkers Prev.* 2002;11:852–861. Allen NE, Appleby PN, Davey GK, et al. The associations of diet with serum insulin-like growth factor I and its main binding proteins in 292 women meat-eaters, vegetarians, and vegans. *Cancer Epidemiol Biomarkers Prev.* 2002;11:1441–1448. Kaklamani VG, Linos A, Kaklamani E, et al. Dietary fat and carbohydrates are independently associated with circulating insulin-like growth factor 1 and insulin-like growth factor-binding protein 3 concentrations in healthy adults. *J Clin Oncol.* 1999;17:3291–3298. Larsson SC, Wolk K, Brismar K, et al. Association of diet with serum insulin-like growth factor I in middle-aged and elderly men. *Am J Clin Nutr.* 2005;81:1163–1167.
 21. Cannata D, Fierz Y, Vijayakumar A, et al. Type 2 diabetes and cancer: what is the connection? *Mt Sinai J Med.* 2010;77:197–213. Venkateswaran V, Haddad AQ, Fleshner NE, et al. Association of diet-induced hyperinsulinemia with accelerated growth of prostate cancer (LNCaP) xenografts. *J Natl Cancer Inst.* 2007;99:1793–1800. Kaaks R. Nutrition, insulin IGF-1 metabolism and cancer risk: a summary of epidemiological evidence. *Novartis Found Symp.* 2004;262:247–260. Discussion 260–68.
 22. Dewell A, Weidner G, Sumner MD, et al. Relationship of dietary protein and soy isoflavones to serum IGF-1 and IGF binding protein in the Prostate Cancer Lifestyle Trial. *Nutr Cancer.* 2007;58(1):35–42. Gann PH, Kazer R, Chatterton R, et al. Sequential, randomized trial of a low-fat, high fiber diet and soy supplementation effects on circulating IGF-1 and its binding proteins in premenopausal women. *Int J Cancer.* 2005;116(2):297–303. Khalil DA, Lucas EA, Juma S, et al. Soy protein supplementation increases serum insulin-like growth factor-1 in young and old men but does not affect markers of bone metabolism *J Nutr.* 2002;132(9):2605–2608.
 23. Sanderson M, Shu XO, Yu H, et al. Insulin-like growth factor-1, soy protein intake, and breast cancer risk. *Nutr Cancer.* 2004;50(1):8–15. Takata Y, Maskarinec G, Rinaldi S, et al. Serum insulin-like growth factor-1 levels among women in Hawaii and Japan with different levels of tofu intake. *Nutr Cancer.* 2006;56(2):136–142.

24. Salvioli S, Capri M, Bucci L, et al. Why do centenarians escape or postpone cancer? The role of IGF-1, inflammation and p53. *Cancer Immunol Immunother.* 2009;58(12):1909–1917.
25. Crowe FL, Key TJ, Allen NE, et al. The association between diet and serum concentrations of IGF-I, IGFBP-1, IGFBP-2, and IGFBP-3 in the European Prospective Investigation into Cancer and Nutrition. *Cancer Epidemiol.* 2009;18(5):1333–1340. Tsilidis KK, Travis RC, Appleby PN, et al. Insulin-like growth factor pathway genes and blood concentrations, dietary protein and risk of prostate cancer in the NCI Breast and Prostate Cancer Cohort Consortium (BPC3). *Int J Cancer.* 2013;133(2):495–504.
26. Fontana L, Weiss EP, Holloszy JO. Long-term effects of calorie or protein restriction on serum IGF-1 and IGFBP-3 concentration in humans. *Aging Cell.* 2008;7(5):681–687.
27. Crowe FL, Key TJ, Allen NE, et al. The association between diet and serum concentrations of IGF-I, IGFBP-1, IGFBP-2, and IGFBP-3 in the European Prospective Investigation into Cancer and Nutrition. *Cancer Epidemiol.* 2009;18(5):1333–1340.
28. Fontana L, Klein S, Holloszy JO. Long-term low-protein, low-calorie diet and endurance exercise modulate metabolic factors associated with cancer risk. *Am J Clin Nutr.* 2006;84:1456–1462.
29. Sinha R, Cross AJ, Graubard BI, et al. Meat intake and mortality: a prospective study of over half a million people. *Arch Intern Med.* 2009;169:562–571. Bernstein AM, Sun Q, Hu FB, et al. Major dietary protein sources and risk of coronary heart disease in women. *Circulation.* 2010;122:876–883. Pan A, Sun Q, Bernstein AM, et al. Red meat consumption and mortality: results from 2 prospective cohort studies. *Arch Intern Med.* 2012;172(7):555–663. Ascherio A, Willett WC, Rimm EB, et al. Dietary iron intake and risk of coronary disease among men. *Circulation.* 1994;89:969–974. Larsson SC, Virtamo J, Wolk A. Red meat consumption and risk of stroke in Swedish men. *Am J Clin Nutr.* 2011;94(2):417–421.
30. Tholstrup T, Hjerpsted J, Raff M: Palm olein increases plasma cholesterol moderately compared with olive oil in healthy individuals. *Am J Clin Nutr.* 2011;94:1426–1432. De Oliveira Otto MC, Alonso A, Lee DH, et al. Dietary intakes of zinc and heme iron from red meat, but not from other sources, are associated with greater risk of metabolic syndrome and cardiovascular disease. *J Nutr.* 2012;142:526–533. Ahluwalia N, Genoux A, Ferrieres J, et al. Iron status is associated with carotid atherosclerotic plaques in middle-aged adults. *J Nutr.* 2010;140:812–816. Brewer GJ. Iron and copper toxicity in diseases of aging, particularly atherosclerosis and Alzheimer's disease. *Exp Biol Med.* 2007;232:323–335.
31. Luan de C, Li H, Li SJ, et al. Body iron stores and dietary iron intake in relation to diabetes in adults in North China. *Diabetes Care.* 2008;31:285–286.

- Rajpathak SN, Crandall JP, Wylie-Rosett J, et al. The role of iron in type 2 diabetes in humans. *Biochim Biophys Acta*. 2009;1790:671–681.
32. Chen GC, Lv DB, Pang Z, et al. Red and processed meat consumption and risk of stroke: a meta-analysis of prospective cohort studies. *Eur J Clin Nutr*. 2013;67:91–95.
33. Kiechl S, Willeit J, Egger G, et al. Body iron stores and the risk of carotid atherosclerosis: prospective results from the Bruneck study. *Circulation*. 1997;96:3300–3307. Steffen LM, Kroenke CH, Yu X, et al. Associations of plant food, dairy product, and meat intakes with 15-y incidence of elevated blood pressure in young black and white adults: the Coronary Artery Risk Development in Young Adults (CARDIA) Study. *Am J Clin Nutr*. 2005;82:1169–1177; quiz 1363–1364. Tzoulaki I, Brown IJ, Chan Q, et al. Relation of iron and red meat intake to blood pressure: cross sectional epidemiological study. *BMJ*. 2008;337:a258. Wang L, Manson JE, Buring JE, et al. Meat intake and the risk of hypertension in middle-aged and older women. *J Hypertens*. 2008;26:215–222. Miura K, Greenland P, Stamler J, et al. Relation of vegetable, fruit, and meat intake to 7-year blood pressure change in middle-aged men: the Chicago Western Electric Study. *Am J Epidemiol*. 2004;159:572–580.
34. Neish AS. Microbes in gastrointestinal health and disease. *Gastroenterology*. 2009;136:65–80. Backhed F. Host responses to the human microbiome. *Nutr Rev*. 2012;70(suppl 1):S14–17.
35. Koeth RA, Wang Z, Levison BS, et al. Intestinal microbiota metabolism of L-carnitine, a nutrient in red meat, promotes atherosclerosis. *Nat Med*. 2013;19(5):576–585. Wang Z, Klipfell E, Bennett BJ, et al. Gut flora metabolism of phosphatidylcholine promotes cardiovascular disease. *Nature*. 2011;472:57–63. Woolston C. Red meat + wrong bacteria = bad news for hearts. *Nature*. <http://www.nature.com/news/red-meat-wrong-bacteria-bad-news-for-hearts-1.12746>. Published April 7, 2013. Accessed April 12, 2013.
36. World Cancer Research Fund, American Institute for Cancer Research. Expert report. Food, nutrition, physical activity and the prevention of cancer: a global perspective. World Cancer Research Fund; 2007. Lunn JC, Kuhnle G, Mai V, et al. The effect of haem in red and processed meat on the endogenous formation of N-nitroso compounds in the upper gastrointestinal tract. *Carcinogenesis*. 2007;28:685–690. Kuhnle GG, Story GW, Reda T, et al. Diet-induced endogenous formation of nitroso compounds in the GI tract. *Free Radic Biol Med*. 2007;43:1040–1047. Pan A, Sun Q, Bernstein AM, et al. Red meat consumption and mortality: results from 2 prospective cohort studies. *Arch Intern Med*. 2012;172(7):555–663. Sinha R, Cross AJ, Graubard BI, et al. Meat intake and mortality: a prospective study of over half a million people. *Arch Intern Med*. 2009;169:562–571. Major JM, Cross AJ, Doubeni CA, et al. Socioeconomic deprivation impact on meat intake and mortality: NIH-AARP Diet and Health Study. *Cancer Causes Control*. 2011;22:1699–1707. Key TJ, Fraser GE, Thorogood M, et al. Mortality in

- vegetarians and nonvegetarians: detailed findings from a collaborative analysis of 5 prospective studies. *Am J Clin Nutr.* 1999;70:516S–524S. Fraser GE. Associations between diet and cancer, ischemic heart disease, and all-cause mortality in non-Hispanic white California Seventh-day Adventists. *Am J Clin Nutr.* 1999;70:532S–538S.
37. Djousse L, Gaziano JM. Egg consumption in relation to cardiovascular disease and mortality: the Physicians' Health Study. *Am J Clin Nutr.* 2008;87:964–969.
 38. Djousse L, Gaziano JM, Buring JE, et al. Egg consumption and risk of type 2 diabetes in men and women. *Diabetes Care.* 2009;32:295–300.
 39. Trichopoulou A, Psaltopoulou T, Orfanos P, et al. Diet and physical activity in relation to overall mortality amongst adult diabetics in a general population cohort. *J Intern Med.* 2006;259:583–591.
 40. Johansson M, Van Guelpen B, Vollset SE, et al. One-carbon metabolism and prostate cancer risk: prospective investigation of seven circulating B vitamins and metabolites. *Cancer Epidemiol Biomarkers Prev.* 2009;18:1538–1543. Platz EA, Clinton SK, Giovannucci E. Association between plasma cholesterol and prostate cancer in the PSA era. *Int J Cancer.* 2008;123:1693–1698.
 41. Richman EL, Kenfield SA, Stampfer MJ, et al. Egg, red meat, and poultry intake and risk of lethal prostate cancer in the prostate specific antigen-era: incidence and survival. *Cancer Prev Res.* 2011;4(12):2110–2121.
 42. Richman EL, Kenfield SA, Stampfer MJ, et al. Choline intake and risk of lethal prostate cancer: incidence and survival. *Am J Clin Nutr.* 2012;96:855–863.
 43. Zeisel, SH. Choline and brain development. In: Bowman BA, Russell RM, eds. *Present Knowledge in Nutrition.* 9th ed. Washington, DC: International Life Sciences Institute; 2006:352–360. Glunde K, Ackerstaff E, Mori N, et al. Choline phospholipid metabolism in cancer: consequences for molecular pharmaceutical interventions. *Mol Pharm.* 2006;3:496–506. Glunde K, Bhujwalla ZM. Choline kinase alpha in cancer prognosis and treatment. *Lancet Oncol.* 2007;8:855–857.
 44. Cho E, Willett WC, Colditz GA, et al. Dietary choline and betaine and the risk of distal colorectal adenoma in women. *J Natl Cancer Inst.* 2007;99:1224–1231.
 45. Spence JD, Jenkins DJ, Davignon J. Egg yolk consumption and carotid plaque. *Atherosclerosis.* 2012;224(2):469–473.

Шеста глава: Планът

1. Streppel MT, Arends LR, Veer P, et al: Dietary fiber and blood pressure: a meta-analysis of randomized placebo-controlled trials. *Arch Intern Med*

- 2005;165:150–156. Lee YP, Puddey IB, Hodgson JM. Protein, Fibre and Blood Pressure: Potential Benefit of Legumes. *Clinical and Experimental Pharmacology and Physiology* 2008; 35(4): 473–476. Papanikolaou Y, Fulgoni VL. Bean Consumption is associated with greater nutrient intake, reduced systolic blood pressure, lower body weight and a smaller waist circumference in adults: Results from the National Health and Nutrition Examination Survey.1999–2002. *J Am Coll Nutr* 2008; 27(5): 569–76.
2. Sievenpiper JL, Kendall CW, Esfahani A, et al. Effect of non-oil-seed pulses on glycaemic control: a systematic review and meta-analysis of randomized controlled experimental trials in people with and without diabetes. *Diabetologia*. 2009;52:1479–1495. Patel A, MacMahon S, Chalmers J, et al. Intensive blood glucose control and vascular outcomes in patients with type 2 diabetes. *N Engl J Med*. 2008;358:2560–2572. Jenkins DA, Kendall CC, Augustin LA, et al. Effect of legumes as part of a low glycemic index diet on glycemic control and cardiovascular risk factors in type 2 diabetes mellitus: a randomized controlled trial. *Arch Intern Med*. 2012;172(21):1653–1660.
 3. Bazzano LA, Thompson AM, Tees MT, et al. Non-soy legume consumption lowers cholesterol levels: a meta-analysis of randomized controlled trials. *Nutr Metab Cardiovasc Dis*. 2011;21:94–103.
 4. Bednar GE, Patil AR, Murray SM, et al. Starch and fiber fractions in selected food and feed ingredients affect their small intestinal digestibility and fermentability and their large bowel fermentability in vitro in a canine model. *J Nutr*. 2001;131:276–286. Muir JG, O’Dea K. Measurement of resistant starch: factors affecting the amount of starch escaping digestion in vitro. *Am J Clin Nutr*. 1992;56:123–127.
 5. Blackberry I, Kouris-Blazos A, Wahlqvist ML, et al. Legumes: the most important dietary predictor of survival in older people of different ethnicities. *Asia Pac J Clin Nutr*. 2004;13(suppl):S126.
 6. Weber Lozada K, Keri RA. Bisphenol A increases mammary cancer risk in two distinct mouse models of breast cancer. *Biol Reprod*. 2011;85(3):490–497. Ho S, Tang W, Prins GS, et al. Developmental exposure to estradio and bisphenol-A increases susceptibility to prostate carcinogenesis and epigenetically regulates phosphodiesterase type 4 variant 4. *Cancer Res*. 2006;66:5624–5632. Melzer D, Rice NE, Lewis C, et al. Association of urinary bisphenol A concentration with heart disease: evidence from NHANES 2003/06. *PLoS One*. 2010;5(1):e8673. Lang IA, Galloway TS, Scarlett A, et al. Association of urinary bisphenol A concentration with medical disorders and laboratory abnormalities in adults. *JAMA*. 2008;300(11):1303–1310. Meeker JD, Ehrlich S, Toth TL, et al. Semen quality and sperm DNA damage in relation to urinary bisphenol A among men from an infertility clinic. *Reprod Toxicol*. 2010;30(4):532–539. Brieno-Enriquez MA, Toran N, Martinez F, et al. Gene expression is altered after bisphenol A exposure in human fetal oocytes in vitro. *Mol Hum Reprod*. 2012;18(4):171–183.

Sugiura-Ogasawara M, Ozaki Y, Sonta S, Makino T, Suzumori K. Exposure to bisphenol A is associated with recurrent miscarriage. *Human Reprod.* 2005;20:2325–2329. Friedrich MJ. Bisphenol A and reproduction. *JAMA.* 2011;305(1):28. Howdeshell KL, Hotchkiss AK, Thayer KA, et al. Environmental toxins: exposure to bisphenol A advances puberty. *Nature.* 1999;401:763–764. Kiguchi M, Fujita S, Oki H, Shimizu N, Cools AR, Koshikawa N. Behavioural characterisation of rats exposed neonatally to bisphenol-A: responses to a novel environment and to methylphenidate challenge in a putative model of attention-deficit hyperactivity disorder. *J Neural Trans.* 2008;115(7):1079–1085. Nagel SC, Boechler M, Welshons WV, et al. Relative binding affinity-serum modified access (RBA-SMA) assay predicts the relative in vivo bioactivity of the xenoestrogens bisphenol A and octylphenol. *Environ Health Perspect.* 1997;105(1):70–76. Li D, Zhou Z, Qing Y, et al. Occupational exposure to bisphenol-A (BPA) and the risk of self-reported male sexual dysfunction. *Human Reprod.* 2010;25(2):519–527. Howdeshell KL, Andrew KH, Thayer KA, et al. Environmental toxins: Exposure to bisphenol A advances puberty. *Nature.* 1999;401:763–764. Braun JM, Yolton K, Dietrich KN, et al. Prenatal bisphenol A exposure and early childhood behavior. *Environ Health Perspect.* 2009;117:1945–1952. Carwile JL, Michels KB. Urinary bisphenol A and obesity: NHANES 2003–2006. *Environ Res.* 2011;111(6):825–830. Wang T, Main L, Bing C, et al. Urinary bisphenol a (BPA) concentration associates with obesity and insulin resistance. *J Clin Endocrinol Metab.* 2012;97(2):E223–227. Bhandari R, Xiao J, Shankar A. Urinary bisphenol A and obesity in U.S. children. *Am J Epidemiol.* 2013;177(11):1263–1270.

Приложение: Да помислим за хранителни добавки

1. Pitkin RM. Folate and neural tube defects. *Am J Clin Nutr.* 2007;85:285S–288S.
2. Troen AM, Mitchell B, Sorensen B, et al. Unmetabolized folic acid in plasma is associated with reduced natural killer cell cytotoxicity among postmenopausal women. *J Nutr.* 2006;136:189–194. Smith AD, Kim YI, Refsum H. Is folic acid good for everyone? *Am J Clin Nutr.* 2008;87:517–533. Pfeiffer CM, Caudill SP, Gunter EW, et al. Biochemical indicators of B vitamin status in the US population after folic acid fortification: results from the National Health and Nutrition Examination Survey 1999–2000. *Am J Clin Nutr.* 2005;82:442–450.
3. Charles D, Ness AR, Campbell D, et al. Taking folate in pregnancy and risk of maternal breast cancer. *BMJ.* 2004;329:1375–1376. Figueiredo JC, Grau MV, Haile RW, et al. Folic acid and risk of prostate cancer: results from a randomized clinical trial. *J Natl Cancer Inst.* 2009;101:432–435. Fife J, Raniga S, Hider PN, et al. Folic acid supplementation and colorectal cancer risk: a meta-analysis. *Colorectal Dis.* 2011;13(2):132–137.
4. Stolzenberg-Solomon RZ, Chang SC, Leitzmann MF, et al. Folate intake, alcohol use, and postmenopausal breast cancer risk in the Prostate,

- Lung, Colorectal, and Ovarian Cancer Screening Trial. *Am J Clin Nutr.* 2006;83:895–904.
5. Baggott JE, Oster RA, Tamura T. Meta-analysis of cancer risk in folic acid supplementation trials. *Cancer Epidemiol.* 2012;36(1):78–81.
 6. Figueiredo JC, Grau MV, Haile RW, et al. Folic acid and risk of prostate cancer: results from a randomized clinical trial. *J Natl Cancer Inst.* 2009;101:432–435. Sellers TA, Kushi LH, Cerhan JR, et al. Dietary folate intake, alcohol, and risk of breast cancer in a prospective study of postmenopausal women. *Epidemiology.* 2001;12:420–428. Martinez ME, Giovannucci E, Jiang R, et al. Folate fortification, plasma folate, homocysteine and colorectal adenoma recurrence. *Int J Cancer.* 2006;119:1440–1446.
 7. Smith AD, Kim YI, Refsum H. Is folic acid good for everyone? *Am J Clin Nutr.* 2008;87:517–533.
 8. Whitrow MJ, Moore VM, Rumbold AR, et al. Effect of supplemental folic acid in pregnancy on childhood asthma: a prospective birth cohort study. *Am J Epidemiol.* 2009;170:1486–1493. Haberg SE, London SJ, Stigum H, et al. Folic acid supplements in pregnancy and early childhood respiratory health. *Arch Dis Child.* 2009;94:180–184. Kwan ML, Jensen CD, Block G, et al. Maternal diet and risk of childhood acute lymphoblastic leukemia. *Public Health Rep.* 2009;124:503–514. Petridou E, Ntouvelis E, Dessypris N, et al. Maternal diet and acute lymphoblastic leukemia in young children. *Cancer Epidemiol Biomarkers Prev.* 2005;14:1935–1939. Jensen CD, Block G, Buffler P, et al. Maternal dietary risk factors in childhood acute lymphoblastic leukemia (United States). *Cancer Causes Control.* 2004;15:559–570. Tower RL, Spector LG. The epidemiology of childhood leukemia with a focus on birth weight and diet. *Crit Rev Clin Lab Sci.* 2007;44:203–242. KalLEN B. Congenital malformations in infants whose mothers reported the use of folic acid in early pregnancy in Sweden: a prospective population study. *Congenit Anom.* 2007;47:119–124.
 9. Higdon J. Carotenoids. In: *An Evidence-Based Approach to Dietary Phytochemicals.* New York: Thieme; 2006: 47–61. Krinsky NI, Johnson EJ. Carotenoid actions and their relation to health and disease. *Mol Aspects Med.* 2005;26:459–516.
 10. The Alpha-Tocopherol Beta Carotene Cancer Prevention Study Group. The effect of vitamin E and beta carotene on the incidence of lung cancer and other cancers in male smokers. *N Engl J Med.* 1994;330:1029–1035. Omenn GS, Goodman GE, Thornquist MD, et al. Effects of a combination of beta carotene and vitamin A on lung cancer and cardiovascular disease. *N Engl J Med.* 1996;334:1150–1155.
 11. Bjelakovic G, Nikolova D, Gluud LL, et al. Antioxidant supplements for prevention of mortality in healthy participants and patients with various diseases. *Cochrane Database Syst Rev.* 2012;3:CD007176.

12. Mayne ST. Beta-carotene, carotenoids, and disease prevention in humans. *FASEB J*. 1996;10:690–701.
13. Krinsky NI, Johnson EJ. Carotenoid actions and their relation to health and disease. *Mol Aspects Med*. 2005;26:459–516.
14. Melhus H, Michaelsson K, Kindmark A, et al. Excessive dietary intake of vitamin A is associated with reduced bone mineral density and increased risk for hip fracture. *Ann Intern Med*. 1998;129:770–778.
15. Higdon J. Vitamin A. In: *An Evidence-Based Approach to Vitamins and Minerals*. New York: Thieme; 2003; 39–47. Botto LD, Loffredo C, Scanlon KS, et al. Vitamin A and cardiac outflow tract defects. *Epidemiology*. 2001;12:491–496.
16. Bjelakovic G, Nikolova D, Gluud LL, et al. Antioxidant supplements for prevention of mortality in healthy participants and patients with various diseases. *Cochrane Database Syst Rev*. 2012;3:CD007176.
17. Vinceti M, Wei ET, Malagoli C, et al. Adverse health effects of selenium in humans. *Rev Environ Health*. 2001;16:233–251. Mueller AS, Mueller K, Wolf NM, et al. Selenium and diabetes: an enigma? *Free Radic Res*. 2009;43:1029–1059. Navas-Acien A, Bley J, Guallar E. Selenium intake and cardiovascular risk: what is new? *Curr Opin Lipidol*. 2008;19:43–49. Chan JM, Oh WK, Xie W, et al. Plasma selenium, manganese superoxide dismutase, and intermediate- or high-risk prostate cancer. *J Clin Oncol*. 2009;27:3577–3583.
18. Brewer GJ. Iron and copper toxicity in diseases of aging, particularly atherosclerosis and Alzheimer's disease. *Exp Biol Med*. 2007;232(2):323–335.
19. Morris MC, Evans DA, Tangney CC, et al. Dietary copper and high saturated and trans fat intakes associated with cognitive decline. *Arch Neurol*. 2006;63(8):1085–1088.
20. Office of Dietary Supplements, National Institutes of Health. Dietary supplement fact sheet: vitamin B12. <http://ods.od.nih.gov/factsheets/VitaminB12/>. Reviewed June 24, 2011.
21. Hooshmand B, Solomon A, Kareholt I, et al. Homocysteine and holotranscobalamin and the risk of Alzheimer disease: a longitudinal study. *Neurology*. 2010;75:1408–1414. Allen LH. How common is vitamin B-12 deficiency? *Am J Clin Nutr*. 2009;89:693S–696S.
22. Hyppönen E. Vitamin D and increasing incidence of type 1 diabetes—evidence for an association? *Diabetes Obes Metab*. 2010;12(9):737–743.
23. Holick MF. Sunlight and vitamin D for bone health and prevention of autoimmune diseases, cancers, and cardiovascular disease. *Am J Clin Nutr*. 2004;80:1678S–1688S. Holick MF, Chen TC. Vitamin D deficiency: a worldwide problem with health consequences. *Am J Clin Nutr*. 2008;87:1080S–1086S.

24. Melamed ML, Michos ED, Post W, et al. 25-Hydroxyvitamin D levels and the risk of mortality in the general population. *Arch Intern Med.* 2008;168:1629–1637. Schöttker B, Haug U, Schomburg L, et al. Strong associations of 25-hydroxyvitamin D concentrations with all-cause, cardiovascular, cancer, and respiratory disease mortality in a large cohort study. *Am J Clin Nutr.* 2013;97(4):782–793. Semba RD, Houston DK, Bandinelli S, et al. Relationship of 25-hydroxyvitamin D with all-cause and cardiovascular disease mortality in older community-dwelling adults. *Eur J Clin Nutr.* 2010;64:203–209. Zittermann A, Iodice S, Pilz S, et al. Vitamin D deficiency and mortality risk in the general population: a meta-analysis of prospective cohort studies. *Am J Clin Nutr.* 2012;95:91–100.
25. Gandini S, Boniol M, Haukka J, et al. Meta-analysis of observational studies of serum 25-hydroxyvitamin D levels and colorectal, breast and prostate cancer and colorectal adenoma. *Int J Cancer.* 2011;128:1414–1424. Grant WB. Relation between prediagnostic serum 25-hydroxyvitamin D level and incidence of breast, colorectal, and other cancers. *J Photochem Photobiol B.* 2010;101:130–136.
26. Tomson J, Emberson J, Hill M, et al. Vitamin D and risk of death from vascular and non-vascular causes in the Whitehall study and meta-analyses of 12,000 deaths. *Eur Heart J.* 2013;34:1365–1374.
27. Hunt JR. Bioavailability of iron, zinc, and other trace minerals from vegetarian diets. *Am J Clin Nutr.* 2003;78(suppl):633S–639S.
28. de Bortoli MC, Cozzolino SM. Zinc and selenium nutritional status in vegetarians. *Biol Trace Elem Res.* 2009;127(3):228–233. Frassinetti S, Bronzetti G, Caltavuturo L, et al. The role of zinc in life: a review. *J Environ Pathol Toxicol Oncol.* 2006;25(3):597–610. Office of Dietary Supplements, National Institutes of Health. Dietary supplement fact sheet: zinc. <http://ods.od.nih.gov/factsheets/Zinc/>. Reviewed June 5, 2013.
29. Gonzalez A, Peters U, Lampe JW, White E. Zinc intake from supplements and diet and prostate cancer. *Nutr Cancer.* 2009;61(2):206–215. Whelan P, Walker BE, Kelleher J. Zinc, vitamin A and prostatic cancer. *Br J Urol.* 1983;55:525–528. Ogunlewe JO, Osegbe DN. Zinc and cadmium concentrations in indigenous blacks with normal, hypertrophic, and malignant prostate. *Cancer.* 1989;63:1388–1392. Feustel A, Wennrich R, Schmidt B. Serum-Zn-levels in prostatic cancer. *Urol Res.* 1989;17:41–42.
30. Waldmann A, Koschizke JW, Leitzmann C, Hahn A. Dietary intakes and lifestyle factors of a vegan population in Germany: results from the German Vegan Study. *Eur J Clin Nutr.* 2003;57:947–955. Krajcovicová-Kudláčková M, Bucková K, Klimes I, Sebková E. Iodine deficiency in vegetarians and vegans. *Ann Nutr Metab.* 2003;47(5):183–185.

Краят на диетите
д-р Джоуел Фърмън

Всички права запазени.
Издавателство Еуниката
София, 2016

www.eunicata.com
facebook.com/Eunicata